

**GAPAC BURNDOWN DEMO - 2003**

**Consultant:** Paul Brown

**Spray Date:** Friday, April 4, 2003

**Location:** Undadilla, GA

No.	Sponsor	Products - Rate Per Acre	1st Rating Date: 4/11/2003					WEED INJURY ON 0 - 5 SCALE*					2nd Rating Date: 4/24/03				
			(A)	(B)	(C)	(D)	(E)	(A)	(B)	(C)	(D)	(E)	(A)	(B)	(C)	(D)	(E)
1.	<a href="#">Valent USA</a>	Valor @ 2.00 OZ. (DRY WT.) + Roundup WeatherMax @ 22 OZ. + Crop Oil @ 16 OZ.	0	1	3	3	2	4.9	4.8	5	5	4.7					
2.	<a href="#">Monsanto</a>	Roundup WeatherMax @ 22 OZ.	0	0	0	3	0	4.7	3	5	5	5					
3.	<a href="#">Syngenta</a>	Gramoxone Max @ 24 OZ. + Caparol @ 24 OZ. + 80/20 Surfactant @ 0.25% v/v	4	1	4	4	4	5	4.5	5	5	5					
4.	<a href="#">Syngenta</a>	Touchdown IQ @ 32.00 OZ. + 2,4-D Amine @ 16.00 OZ.	2	3	2	3	.-	5	4.9	5	5	.-					
5.	<a href="#">CJB</a>	Liberty @ 16 OZ. + Scorch @ 16 OZ. (DRY WT.)	1	1	1	3	2	1	0	4	2	2					
6.	<a href="#">Bayer</a>	Liberty @ 16 OZ. + Roundup WeatherMax @ 16 OZ. + AMS @ 48 OZ. (DRY WT.)	2	2	2	3	2	4.9	4.9	5	5	3					
7.	<a href="#">Bayer</a>	Liberty @ 32 OZ. + AMS @ 48 OZ. (DRY WT.)	2	3	3	3	2	5	5	5	5	5					
8.	Untreated Check		0	0	0	0	0	0	0	0	0	0					
9.	<a href="#">FMC</a>	Aim @ 1.00 OZ. + Diuron @ 10 OZ. (DRY WT.) + Crop Oil @ 1.00 % v/v	2	1	2	2	0	3.5	1	2	4	0					
10.	<a href="#">CJB</a>	Liberty @ 16 OZ. + Scorch @ 8 OZ. (DRY WT.)	2	2	1	3	1	4	1	2	3	2					
11.	<a href="#">CJB</a>	Liberty @ 16 OZ. + Scorch @ 16 OZ. (DRY WT.)	0	2	2	3	0	4	3.5	4	4.5	3.5					
12.	<a href="#">GAPAC Standard</a>	Roundup WeatherMax @ 22.00 OZ. + 2,4-D Amine @ 16 OZ.	2.5	2.5	3	3	1	5	4.5	4	5	4					
13.	<a href="#">GAPAC Standard</a>	Gramoxone Max @ 28 OZ. + Diuron @ 10 OZ. (DRY WT.)	4	1	5	.-	5	5	4.5	5	.-	5					

\*0 = No Injury 1 = Light injury. Burned leaf edges. Slight wilt. 2 = Light/moderate injury. 3 = Moderate injury. 4 = Highly injured. Plants brown - green stem interior. 5 = Dead plants.

Key	Weed Identity	Density per sq ft	Growth Stage
(A) =	C. Geranium		
(B) =	CLE Primrose		
(C) =	Curly Dock		
(D) =	Grasses		
(E) =	False Dandelion		

COMMENTS 4/11/2003: Plots are heavily infested with Carolina geranium, uniform medium infestation of cutleaf evening primrose. Curly dock and false dandelion are scattered throughout the plots. Grasses are primarily identified as blue grass.

